8900271

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Virginia Polytechnic Institute and State University

Williereas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-LUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT Y THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT.

United States seed of this variety (1) shall be sold by variety name only as OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'Camp'

In Testimony Wancroot, I have hereunto set my hand and caused the seal of the Blant Taxiety Protection Office to be affixed at the City of Washington, D. C. this 30th day of November the year of our Lord one thousand nine hundred and eighty-nine.

Plant Variety Protection Office

| U.S. DEPARTM AGRICULTURA APPLICATION FOR PLANT VA (Instruc | FORM APPROVED: OMB NO. 0581-0066 Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426). | | | | | |
|---|---|--|---------------------|---|--|--|
| 1. NAME OF APPLICANT(S) Va. Polytechnic Institute & State | University | 2. TEMPORARY DESIGNATION V81–1325–62 | 1 | 3. VARIETY NAME Camp | | |
| 4. ADDRESS (Street and No. or R.F.D. No., City, | State, and Zip Code) | 5. PHONE (Include area code) | ╁┈ | FOR OFFICIAL USE ONLY | | |
| Blacksburg, Virginia 24061 | t in the second | 703/231–6483 | PVP | 8900271 | | |
| 6. GENUS AND SPECIES NAME | 7. FAMILY NAI | ME (Botanical) | | DATE July 12 1989 | | |
| Glycine max | Leguminosa | e | FILING | TIME 10:00 [Xa.M.]P.M. | | |
| 8. KIND NAME | 9. | DATE OF DETERMINATION | 1 | AMOUNT FOR FILING | | |
| Soybeans | | November 1, 1988 | RECEIVED | \$2150 DAME July 121989 | | |
| 10. IF THE APPLICANT NAMED IS NOT A "PE- partnership, association, etc.) University | RSON," GIVE FORM | OF ORGANIZATION (Corporation | FEES RE | \$ 250. | | |
| | rantin di seriesa di s | | | Nov.1,1989 | | |
| 11. IF INCORPORATED, GIVE STATE OF INCO | PROPATION | | 12. | DATE OF INCORPORATION | | |
| a. | ariety (Request form Variety. Applicant's Ownership SEEO OF THIS VARII | from Plant Variety Protection Off | ice.) | Y AS A CLASS OF CERTIFIED | | |
| 16. DOES THE APPLICANT(S) SPECIFY THAT LIMITED AS TO NUMBER OF GENERATION | THIS VARIETY BE | 17. IF "YES" TO ITEM 16, | WHICH | CLASSES OF PRODUCTION | | |
| X Yes No | | BEYOND BREEDER SE | | egistered Certified | | |
| 18. DID THE APPLICANT(S) PREVIOUSLY F | | ON OF THE VARIETY IN THE L | J.S.? | Yes (If "Yes," give date) No | | |
| Public release in the U.S. was Feb date. No foreign releases have be | oruary 1, 1989, i | | | | | |
| 20. The applicant(s) declare(s) that a viable significant plenished upon request in accordance with | ample of basic seeds h such regulations a | of this variety will be furnishe is may be applicable. | d with | the application and will be re- | | |
| The undersigned applicant(s) is (are) the edistinct, uniform, and stable as required is Variety Protection Act. | owner(s) of this sex n Section 41, and is | ually reproduced novel plant va entitled to protection under th | riety, : ie prov | and believe(s) that the variety is issions of Section 42 of the Plant | | |
| Applicant(s) is (are) informed that false re | epresentation herein | can jeopardize protection and | | | | |
| SIGNATURE OF APPLICANT Scarm A. O | Bun | | 0 | July 10, 1989 | | |
| SIGNATURE OF APPLICANT | | Swiger, Assoc. Dire | ctor | June 30, 1989 | | |

FORM LS-470 (3-86)

Edition of 7-84 obsolete.

Attachment 14a

Application for Plant Variety Protection Certificate for Camp Soybeans

Exhibit A, Origin and Breeding History of Camp Soybeans

Camp is a selection made from the variety Vance which was also developed and released by the Virginia Agricultural Experiment Station. Vance was derived from an F_5 single plant selection from a cross of Essex x $\underline{\text{Glycine}}$ $\underline{\text{soja}}$. The identity of the male parent was lost, but it was typical of the $\underline{\text{Glycine}}$ $\underline{\text{soja}}$ species, possessing a very viny growth habit, small leaves and very small, black seeds. Very intensive selection was conducted in the early generations for plant type similar to good, adapted varieties.

Vance was released as a variety in 1986. That year a non-replicated progeny-row planting was made from 78 individual plants that had been randomly chosen from Vance in 1985 to be used in purification of the variety and for Breeder Seed production. However, some variation among the rows was observed for plant height, leaflet shape, maturity, etc. Forty-nine of these rows were selected as being typical of Vance in general appearance and were harvested individually. Seed size and percent of seeds with green seedcoats were determined for each sub-line. Significant variation among the lines was observed for both traits. Green seedcoats are considered objectionable for natto production, so fourteen of the lines were selected for small seed size and low proportion of seeds with green seedcoats. These were put in a replicated field trial at one location in 1987 and small increase blocks of eight of them were planted. Based on data from this experiment, two sub-lines, V81-1325-10 and V81-1325-62 were very similar and had the least amount of green seedcoats. Since they had promise as improvements over Vance, about 30 lbs. of seed of each was sent to Belize for winter increase. In 1988, the seed returned from Belize was planted for further increase and the two lines, as well as three others, were planted in replicated tests at three locations for further evaluations.

A summary of all available data indicated that V81-1325-62 was most like Vance in yield, maturity, plant height, etc., but had significantly smaller and more uniform seed size as well as a much lower proportion of green seedcoats. The latter properties made it more desirable than Vance for natto production, so a decision was made to release it as a new variety, Camp.

A portion of the 1988 seed increase field of V81-1325-62 was carefully rogued at maturity and harvested to serve as Breeder Seed for future increases.

Attachment 14b

Application for Plant Variety Protection Certificate for Camp Soybeans

Exhibit B. Novelty Statement

Camp is easily distinguished from most other varieties by its small seed size. Since it is a selection from Vance, it is very similar to Vance in general appearance, but has consistently smaller seeds than Vance. In four tests in different environments in 1987 and 1988. Camp seeds ranged from 1.2 gm/100 seeds to 0.7 gm/100 seeds smaller than those of Vance. The average difference was 1.0 gm/100 seeds with 0.3 gm/100 seeds required for statistical significance at the 0.05 probability level. Similarly, the proportion of seeds that passed through a 14/64" round-hole screen averaged 72% for Vance and 90% for Camp with 3% required for statistical significance.

We are aware of a few other small-seeded soybean varieties, but most of them are in earlier maturity groups. The only other variety of Group V maturity which we have been able to compare to Camp is one designated Rocky. It matures about 3 days earlier than Camp, is about 4 inches taller and is much more prone to lodging. The most obvious distinguishing features, however, are the round or ovate leaflets and white flowers of Rocky vs. the lanceolate leaflets and purple flowers of Camp. Hartz Seed Co. has some small-seeded varieties of Group V maturity or later, but these are not generally available for public use and have not been compared to Camp.

EXHIBIT C (Soybean)

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT, GRAIN & SEED DIVISION PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY SOYBEAN (Glycine max L.)

| NAME OF APPLICANT(S) | TEMPORARY DESIGNATION | VARIETY NAME |
|--|--|--|
| Va. Polytechnic Institute & State Univ. | V81-1325-62 | Camp |
| ADDRESS (Street and No., or R.F.D. No., City, State, and Zip Coo | de) | FOR OFFICIAL USE ONLY |
| Blacksburg, Virginia 24061 | | 8900271 |
| Choose the appropriate response which characterizes the va- in your answer is fewer than the number of boxes provided, | riety in the features described , place a zero in the first box w | below. When the number of significant digits then number is 9 or less (e.g., 0 9). |
| 1. SEED SHAPE: | T 2 = Spherical Flattened | (L/W ratio > 1.2; L/T ratio = < 1.2) (L/T ratio > 1.2; T/W > 1.2) |
| 2. SEED COAT COLOR: (Mature Seed) | | |
| 1 = Yellow 2 = Green 3 = Brown | 4 = Black 5 = Other | (Specify) |
| 3. SEED COAT LUSTER: (Mature Hand Shelled Seed) | | |
| 2 1 = Dull ('Corsoy 79'; 'Braxton') 2 = Shiny ('Nebso | oy"; "Gasoy 17") | |
| 4. SEED SIZE: (Mature Seed) | · · · · · · · · · · · · · · · · · · · | |
| 0 8 Grams per 100 seeds | | |
| 5. HILUM COLOR: (Mature Seed) | | |
| 2 1 = Buff 2 = Yellow 3 = Brown | 4 = Gray 5 = Imperfect Blad | ck 6 = Black 7 = Other (Specify) |
| 6. COTYLEDON COLOR: (Mature Seed) | | |
| 1 = Yellow 2 = Green | | |
| 7. SEED PROTEIN PEROXIDASE ACTIVITY: | | |
| 1 = Low 2 = High | | |
| 8. SEED PROTEIN ELECTROPHORETIC BAND: | | |
| 1 = Type A (SP1 ^a) 2 = Type B (SP1 ^b) | | |
| 9. HYPOCOTYL COLOR: | | |
| 1 = Green only ('Evans'; 'Davis') 2 = Green with 3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson'; ' | n bronze band below cotyledons ('V 'Coker Hampton 266A') | Noodworth'; 'Tracy') |
| 0. LEAFLET SHAPE: | | |
| 1 = Lanceolate 2 = Oval 3 = Ovate | 4 = Oshor (0 | |
| 1 = Lanceolate 2 = Oval 3 = Ovate | 4 = Other (Specify) | - // |

FORM LMGS-470-57 (2-82)

| , 11. LEAFLET SIZE: | |
|--|-------------------------------|
| 1 = Smail ('Amsoy 71'; 'A5312') 2 = Medium ('Corsoy 79'; 'Gasoy 17') 3 = Large ('Crawford'; 'Tracy') | |
| 12. LEAF COLOR: | |
| 1 = Light Green ('Weber'; 'York') 2 = Medium Green ('Corsoy 79'; 'Braxto') 3 = Dark Green ('Gnome'; 'Tracy') | on') |
| 13, FLOWER COLOR: | |
| 1 = White 2 = Purple 3 = White with purple throat | |
| 14, POD COLOR: | |
| 1 = Tan 2 = Brown 3 = Błack | |
| 15. PLANT PUBESCENCE COLOR: | |
| 1 = Gray 2 = Brown (Tawny) | |
| 16. PLANT TYPES: | |
| 1 = Slender ('Essex'; 'Amsoy 71') 2 = Intermediate ('Amcor'; 'Braxton') 3 = Bushy ('Gnome'; 'Govan') | |
| 17. PLANT HABIT: | |
| 1 = Determinate ('Gnome'; 'Braxton') 2 = Semi-Determinate ('Will') 3 = Indeterminate ('Nebsoy'; 'Improved Pelican') | |
| | |
| 18. MATURITY GROUP: | |
| 18. MATURITY GROUP: 0 8 | 7 = IV 8 = V |
| 0 8 1 = 000 2 = 00 3 = 0 4 = I 5 = II 6 = III 9 = VI 10 = VII 11 = VIII 12 = IX 13 = X | 7 = IV 8 = V |
| 0 8 1 = 000 2 = 00 3 = 0 4 = I 5 = II 6 = III | 7 = IV 8 = V |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: | 7 = IV 8 = V |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) | 7 = IV 8 = V |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Bacterial Blight (Pseudomonas glycinea) | 7 = IV 8 = V |
| 1 = 000 2 = 00 3 = 0 4 = I 5 = II 6 = III 9 = VI 10 = VII 11 = VIII 12 = IX 13 = X 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) | 7 = IV 8 = V |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: | 7 = IV 8 = V |
| 1 = 000 2 = 00 3 = 0 4 = I 5 = II 6 = III 9 = VI 10 = VII 11 = VIII 12 = IX 13 = X 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) | 7 = IV 8 = V |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) | 7 = IV 8 = V |
| 0 8 1 = 000 2 = 00 3 = 0 4 = I 5 = II 6 = III 9 = VI 10 = VII 11 = VIII 12 = IX 13 = X 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: 0 Bacterial Pustule (Xanthomonas phaseoli var. sojensis) 0 Bacterial Blight (Pseudomonas glycinea) 0 Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: 0 Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) 0 Race 1 Race 2 Race 3 Race 4 Race 5 | 7 = IV 8 = V Other (Specify) |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) | |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2 Race 3 Race 4 Race 5 Downy Mildew (Peronospora trifoliorum var. manshurica) | |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2 Race 3 Race 4 Race 5 | |
| 19. DISEASE REACTION: (Enter 0 = Not Tested; 1 = Susceptible; 2 = Resistant) BACTERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli var. sojensis) Bacterial Blight (Pseudomonas glycines) Wildfire (Pseudomonas tabaci) FUNGAL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 | |

| 19. DISEASE REACTIO | N: (Enter 0 = Not Tested; 1 = Susceptible; 2 = | Resistant) (Continued) | | |
|----------------------|---|------------------------|---------------------------------------|--------|
| FUNGAL DISEA | SES: (Continued) | | | |
| O Pod and St | em Blight <i>(Diaporthe phaseolorum</i> var; sojae) | | | |
| O Purple Seed | l Stain (Cercospora kikuchii) | | | |
| 0 Rhizoctoni | a Root Rot (Rhizoctonia solani) | | | |
| Phytophthe | ora Rot (Phytophthora megasperma var. sojae) | _ | | |
| 0 Race 1 | 0 Race 2 0 Race 3 0 | Race 4 0 Race 5 | Race 6 | Race 7 |
| 0 Race 8 | O Race 9 Other (Specify) | | | |
| VIRAL DISEASES | S: | | | |
| 0 Bud Blight | (Tobacco Ringspot Virus) | • | · · · · · · · · · · · · · · · · · · · | |
| O Yellow Mos | aic (Bean Yellow Mosaic Virus) | · | | |
| O Cowpea Mo | saic (Cowpea Chlorotic Virus) | | | |
| O Pod Mottle | (Bean Pod Mottle Virus) | | | |
| Seed Mottle | (Soybean Mosaic Virus) | | | |
| NEMATODE DISE | ASES: | | | |
| Soybean Cy: | st Nematode (Heterodera glycines) | | | |
| 1 Race 1 | 1 Race 2 1 Race 3 1 | Race 4 Other | (Specify) | |
| 0 Lance Nema | tode (Hopiciaimus Colombus) | | | |
| | ot Knot Nematode (Meloidogyne incognita) | | | |
| | ot Knot Nematode (Meloidogyne Hapla) | | | • |
| | Knot Nematode (Meloidogyne arenaria) | | | |
| | matode (Rotylenchulus reniformis) | | | |
| | EASE NOT ON FORM (Specify): | | | |
| | ENGLINOT ON TOTAL Speciny). | | | |
| 0. PHYSIOLOGICAL RE | SPONSES: (Enter 0 = Not Tested; 1 = Suscept | tible; 2 = Resistant) | | |
| 0 Iron Chlorosi | s on Calcareous Soil | | | |
| Other /Specif | (v) | | | • |
| 1. INSECT REACTION: | (Enter 0 = Not Tested; 1 = Susceptible; 2 = Re | sistant) | | |
| 0 Mexican Bear | Beetle (<i>Epilachna varivestis)</i> | | | |
| O Potato Leaf H | lopper (Empoasca fabae) | | | |
| Other (Specif | y) | | | |
| 2. INDICATE WHICH VA | ARIETY MOST CLOSELY RESEMBLES THA | T SUBMITTED | | |
| CHARACTER | NAME OF VARIETY | CHARACTER | NAME OF VA | RIETY |
| Plant Shape | Vance | Seed Coat Luster | Vance | |
| Leaf Shape | Vance | Seed Size | Vance | |
| Leaf Color | Vance | Seed Shape | Vance | |
| Leaf Size | Vance | Seedling Pigmentation | Vance | |
| | · · | | | |

23. GIVE DATA FOR SUBMITTED AND SIMILAR STANDARD VARIETY: Paired Comparison Data

| | , , , , , , , , , , , , , , , , , , , | | - Carlo | | | | | | |
|-------------------------------------|---|----------|------------|--------------|-----------|--------------|-------|----------------|---------------|
| VARIETY | NO. OF PLANT DAYS LODGIN | | GING PLANT | LEAFLET SIZE | | SEED CONTENT | | SEED SIZE | NO. |
| | MATURITY | SCORE | HEIGHT | CM Width | CM Length | % Protein | % Oil | G/100 SEEDS | SEEDS/ POD |
| Submitted ' | 136 | 1.0 | 60 | | | 39.6 | 19.8 | 7,8 | |
| Vance Name of Similar Variety | 137 | 1.0 | 63 | | | 40.0 | 19.3 | 8.8 | |
| PURI ICATIO | ONS LISSEL | 10.00000 | | <u> </u> | | | | | |

PUBLICATIONS USEFUL AS REFERENCE AIDS FOR COMPLETING THIS FORM:

- 1. Caldwell, B.E., ed. 1973. Soybeans: Improvement, Production, and Uses. Amer. Soc. Agron. Monograph No. 16.
- 2. Buttery, B.R. and R.I. Buzzell. 1968. Peroxidase activity in seeds of soybean varieties. Crop Sci., 8: 722-725.
- 3. Hymowitz, T. 1973. Electrophoretic analysis of SBTI-A2 in the USDA soybean germplasm collection. Crop Sci., 13: 420-421.
- 4. Payne, R.C. and L.F. Morris. 1976. Differentiation of soybean cultivars by seedling pigmentation patterns. J. Seed Technol. 1: 1-19.

Attachments 14c, 14d and 14e

Application for Plant Variety Protection Certificate for Camp

- 14c. Exhibit C attached
- 14d. No additional information necessary.
- 14e. The breeder of this variety is Glenn R. Buss, who is an employee of the applicant, Virginia Polytechnic Institute and State University.